



Globally Ranked Species Richness: G1, G2, G3

These EnviroAtlas national maps display the number of NatureServe Globally Ranked Species (G1, G2, and G3) with potential habitat within each 12-digit hydrologic unit ([HUC](#)) in the conterminous United States. These data are based on habitat models, not wildlife counts.

Why are globally ranked species important?

The NatureServe Globally Ranked Species Richness (G1, G2, and G3) metric estimates the number of species within 3 conservation status tiers based on potential habitat.¹ G1 species are critically imperiled, G2 species are imperiled, and G3 species are vulnerable. NatureServe uses 10 factors within 3 categories—rarity, threats, and trends—to assess conservation status. Each category has 2 to 6 conservation status factors to ensure information used to assign conservation status is consistent and rigorous. Rarity factors include population size, range extent, area of occupancy, number of occurrences, numbers or percent area occupied with good viability and ecological integrity, and environmental specificity. Trend factors include long-term and short-term trends in population size or area. The threat factors are based on the scope and severity of major threats and intrinsic vulnerability.^{1,2}

Species richness is one measure of [biodiversity](#) that can represent the relative conservation value of a particular area. Many scientists believe that biodiversity, because it represents all forms of life on earth, provides or supports the core benefits that humans derive from their environment to sustain human society, economy, health, and well-being. Managing for biodiversity is one way to balance competing demands for ecosystem services.³

Each species plays an important role within its [ecosystem](#), and ecosystems are highly interconnected. Each species depends on others for some aspect of its survival to provide food, habitat, decomposition, pollination, or control of pest species. The removal of even one species from an ecosystem can create a [trophic cascade](#) that can affect the entire [food chain](#). For example, grazers and browsers directly modify the species composition, diversity, and condition of grassland and forest habitats. Top predators, by regulating herbivore numbers, indirectly influence habitat condition and diversity by reducing grazing pressure on plant production.⁴

Characteristics of biodiversity are valued in a variety of ways and thus they are important to include in any assessment that



Photo: Idaho ground squirrel, Dennis Mackey, USFWS

seeks to identify and quantify the value of ecosystems to humans. Some biodiversity metrics clearly reflect ecosystem services and their contribution to our quality of life and economy (e.g., abundance and diversity of game species), whereas others reflect indirect and difficult to quantify relationships to services (e.g., relevance of species diversity to ecosystem resilience, cultural value, and aesthetic amenities).

How can I use this information?

Three EnviroAtlas maps, Mean, Maximum, and Normalized Index of Biodiversity (NIB) illustrate NatureServe Globally Ranked Species Richness (G1, G2, and G3) for each 12-digit HUC in the conterminous United States.⁵ Used together or independently, these maps can help identify areas of low or high potential species richness to help inform decisions about resource restoration, use, and conservation. Mean richness is a commonly used and understood value for comparison. NIB provides an index to compare a metric with other metrics across multiple project scales simultaneously. Maximum richness identifies areas that are species rich but may not occupy large areas (e.g. linear riparian areas).

These maps can be used in conjunction with other maps in EnviroAtlas such as ecoregions, the U.S. Geological Survey (USGS) protected areas database ([PAD-US](#)), or the USGS Gap Analysis Project ([GAP](#)) ecological systems to identify areas with high ecological or recreational value for conservation, recreation, or restoration planning.

After learning the globally ranked species richness values for a particular 12-digit HUC, users can investigate an area more intensively by using individual species models available from the GAP Project.

How were the data for this map created?

The USGS GAP project maps the distribution of natural vegetation communities and potential habitat for individual terrestrial vertebrate species. These models use environmental variables (e.g., land cover, elevation, and distance to water) to predict habitat for each species. GAP modeled habitat for 1,590 terrestrial vertebrate species that reside, breed, or use the habitat within the conterminous U.S. for a significant portion of their life history. This map is based on NatureServe Globally Ranked Species (G1, G2, G3) predicted habitat within the conterminous United States.

Predicted habitat for the 255 individual NatureServe globally ranked species was combined to calculate species richness by pixel. Of the 255 species, there were 105 amphibians, 29 birds, 58 mammals and 63 reptile species. The mean and maximum numbers of globally ranked terrestrial vertebrate species in each 30-meter pixel were calculated for each 12-digit HUC. The mean species richness value by HUC was divided by the maximum mean value within all HUCs to calculate the NIB.

What are the limitations of these data?

EnviroAtlas uses the best data available, but there are still limitations associated with these data. These data, based on models and large national geospatial databases of predicted habitat, are estimations of reality that may overestimate actual species presence. Modeled data are intended to complement rather than replace monitoring data. Habitat models do not predict the actual occurrence of species, but rather their potential occurrence based on their known associations with certain habitat types. Habitat is only one factor that determines

the actual presence of a species. Other factors include habitat quality, predators, prey, competing species, and fine scale habitat features.

Other essential species information in addition to species richness includes the types of species and their [functional groups](#), whether they are rare or common, native or non-native, tolerant or intolerant of disturbance.

How can I access these data?

EnviroAtlas data can be viewed in the interactive map, accessed through web services, or downloaded. Individual 30-meter pixel data may be downloaded from the [New Mexico State University Center for Applied Spatial Ecology](#).

Where can I get more information?

A selection of resources related to terrestrial vertebrates and biodiversity is listed below. Information on the models and data used in the USGS Core Science Analytics, Synthesis & Library's [GAP](#) project is available on their website. For additional information on how the data were created, access the [metadata](#) for the data layer from the layer list drop down menu. To ask specific questions about this data layer, please contact the [EnviroAtlas Team](#).

Acknowledgments

The data for NatureServe Globally Ranked Species Richness were created through a collaborative effort between the EPA and USGS GAP. GAP habitat models were created by New Mexico State University, North Carolina State University, and Boise State University personnel. Kenneth Boykin and graduate students from New Mexico State University generated the biodiversity metrics. The fact sheet was written by Kenneth Boykin (New Mexico State University), William Kepner, Anne Neale, and Jessica Daniel (EPA), and Megan Culler (EPA Student Services Contractor).

Selected Publications

1. NatureServe. 2020. [NatureServe Conservation Status Assessment: Identifying threatened species and ecosystems](#). Accessed April 2020.
2. Faber-Langendoen, D., J. Nichols, L. Master, K. Snow, A. Tomaino, R. Bittman, G. Hammerson, B. Heidel, L. Ramsay, A. Teucher, and B. Young. 2012. [NatureServe conservation status assessments: Methodology for assigning ranks](#). NatureServe, Arlington, VA.
3. Boykin, K.G., W.G. Kepner, D.F. Bradford, R.K. Guy, D.A. Kopp, A. Leimer, E. Samson, F. East, A. Neale, and K. Gergely. 2013. [A national approach for mapping and quantifying habitat-based biodiversity metrics across multiple spatial scales](#). *Ecological Indicators* 33:139–147.
4. Miller, B., B. Dugelby, D. Foreman, C. Martinez del Rio, R. Noss, M. Phillips, R. Reading, M. E. Soulé, J. Terborgh, and L. Wilcox. 2001. [The importance of large carnivores to healthy ecosystems](#). *Endangered Species Update* 18(5):202–210.
5. Kepner, W.G., K.G. Boykin, D.F. Bradford, A.C. Neale, A.K. Leimer, and K.J. Gergely. 2011. [Biodiversity metrics fact sheet](#), EPA/600/F-11/006, U.S. Environmental Protection Agency, Washington, D.C.